For most functions f a proof of  $\lim_{x \to +\infty} f(x) = L$  based on the definition in the notes should consist from the following steps.

- (1) Find  $X_0$  such that f(x) is defined for all  $x \ge X_0$ . Justify your choice.
- (2) Use algebra to simplify the expression |f(x) L| with the assumption that  $x \ge X_0$ . Try to eliminate the absolute value.
- (3) Use the simplification from (2) to discover a BIN:

 $||f(x) - L| \le b(x) \text{ valid for } x \ge X_0$ The content of the box above is a BIN.

Here b(x) should be a simple function with the following properties:

- (a) b(x) > 0 for all  $x \ge X_0$ .
- (b) b(x) is tiny for huge x.

(c)  $b(x) < \epsilon$  is easily solvable for x for each  $\epsilon > 0$ . The solution should be of the form

x > some expression involving  $\epsilon$ 

**Warning:** In the above inequality some expression involving  $\epsilon$  must be huge when  $\epsilon$  is tiny.

(4) Use the solution of  $b(x) < \epsilon$ , that is some expression involving  $\epsilon$ , and  $X_0$  to define

$$X(\epsilon) = \max \left\{ X_0, \text{ some expression involving } \epsilon \right\}$$

(5) Use the BIN above to prove the implication x > X(ε) ⇒ |f(x) − L| < ε.</li>
Note: The structure of this proof is always the same.

- (i) First assume that  $x > X(\epsilon)$ .
- (ii) The definition of  $X(\epsilon)$  yields that

 $X(\epsilon) \ge X_0$  and  $X(\epsilon) \ge$  some expression involving  $\epsilon$ 

(iii) Based of (5i) and (5ii) we conclude that the following two inequalities are true:

 $x > X_0$  and x > some expression involving  $\epsilon$ 

(iv) From (3) part (c) we know that

$$x >$$
some expression involving  $\epsilon$  implies  $b(x) < \epsilon$ .

Therefore (5iii) yields that  $b(x) < \epsilon$  is true.

(v) We also established that the BIN is true:

$$|f(x) - L| \le b(x)$$
 valid for  $x \ge X_0$ 

(vi) Together  $|f(x) - L| \le b(x)$  and  $b(x) < \epsilon$  yield

$$|f(x) - L| < \epsilon.$$

This is exactly what we needed to prove.